

Report for Drucker Labs


Sample ID	Brunswick Lab ID	ORAC _{hydro} * (μ moleTE/L)	ORAC _{lipo} ^ (μ moleTE/L)	ORAC _{total} (μ moleTE/L)
IntraMAX Lot# 415	06-1457	54,982	1,041	56,023

*The ORAC analysis provides a measure of the scavenging capacity of antioxidants against the peroxy radical, which is one of the most common reactive oxygen species (ROS) found in the body. ORAC_{hydro} reflects water-soluble antioxidant capacity and the ^ ORAC_{lipo} is the lipid soluble antioxidant capacity. Trolox, a water-soluble Vitamin E analog, is used as the calibration standard and the ORAC result is expressed as micromole Trolox equivalent (TE) per liter.

The acceptable precision of the ORAC assay is 15% relative standard deviation.¹⁻²

Testing performed by J. Frietas.

Approved by:


Boxin Ou, PhD.
Vice President

B-4349 / 5-10-06 jt

Samples will be discarded one month from report date, unless otherwise notified by customer in writing.

¹ Ou, B; Hampsch-Woodill, M.; Prior, R. L.; Development and Validation of an Improved Oxygen Radical Absorbance Capacity Assay using Fluorescein as the Fluorescent Probe. Journal of Agricultural and Food Chemistry.; **2001**; 49(10); 4619-4626

² Huang, D.; Ou, B.; Hampsch-Woodill, M.; Flanagan, J.; Deemer, E. K.; Development and Validation of Oxygen Radical Absorbance Capacity Assay for Lipophilic Antioxidants using Randomly Methylated β -Cyclodextrin as the Solubility Enhancer. Journal of Agricultural and Food Chemistry.; **2002**, 50(7); 1815-1821.